

Parathyroid Scan (Tc-99m)

Introduction

A parathyroid scan is a nuclear medicine exam done to determine the anatomy and function of your parathyroid glands. This may be to investigate a high blood calcium level.

Preparation

Inform us if you are on any thyroid medication as this may need to be stopped for the scan.

Inform us if you have had any scans involving iodine contrast in the 6 weeks prior to the scan

The injection will not impair your ability to drive a vehicle.

Pregnant women or children should not attend the appointment with you.

Ensure that you are well hydrated.

Women of childbearing age

This scan should be scheduled within 28 days of the commencement of your last menstrual period to avoid inadvertent irradiation of a foetus. If your period is overdue, your appointment may be rearranged.

Scan

On arrival you will be given the isotope injection into a vein in your arm. This injection should have no side effects. You will be asked to drink a glass of water to clear your salivary glands. 20 minutes after the injection, you will be asked to lie on the scanner table and a ten minute image will be taken.

1 hour after the injection, you will be asked to lie on the scanner table again and a ten minute image will be taken.

2 hours after the injection, you will be asked to lie on the scanner table for the final scan. This time, the scanner will move around you. This scan will take approximately 30 minutes.

Patient Information Leaflet

Radiation precautions after the scan

Some of the isotope will remain in your body for approximately 24 hours after your scan is complete. To minimise the radiation dose to yourself and others, you should take the following precautions during this period:

1. Drink plenty of fluids and empty your bladder as frequently as possible. When using the toilet you should avoid spills, flush the toilet twice and wash your hands thoroughly. You should sit on the toilet when passing urine.
2. If possible, avoid close contact with pregnant women, infants and children for 24 hours after your injection.
3. If you are a nursing mother express and discard breast milk for 24 hours after your injection. You may resume normal breast-feeding after this time.
4. You should postpone any other medical tests or procedures for 24 hours after the injection where possible.

Medical radiation: risks v benefits

We are all exposed to natural background radiation every day. Medical exposures give a small additional dose on top of natural radiation.

The amount of radiation received during a nuclear medicine procedure is low, resulting in the equivalent of approximately a few months to two year's background radiation.

The only effect on the patient that is known to be possible at these low doses is a very slight increase in the chance of cancer occurring many years or decades after the exposure.

As long as it is clearly necessary to help make the correct diagnosis and treatment decision, the benefits of detection, diagnosis and treatment resulting from the nuclear medicine examination should outweigh these small radiation risks.